

Nonstationary boundary value problems for wave equation and their generalized solutions

Alexeyeva L.A.

*Institute of Mathematics, Kazakhstan
Pushkin str.125, Almaty, 050010, alexeeva@math.kz*

Investigation of many problems of acoustics, hydromechanics, elastodynamics and others is connected with boundary value problem (BVP) for wave equation - the multidimensional analogue of the D'Alambert's equation, which describes processes of spreading the waves in uniform isotropic ambience. So elaboration of the efficient methods of their solving for areas with free geometry and varied type of the border conditions is very currently .

The most efficient method of the study of such problems is the boundary integral equations method (BIEM). Its main advance is reduction of dimension of solved equations and increasing calculation stability. That is very greatly, particularly for unbounded areas. The existing methods and programs of splines approximation for free contours and surfaces remove the problem of restriction of the form of considered areas by using BIEM. At present time this method is broadly used for solving elliptical and parabolic problems, what is connected with success in development of BIE theories for elliptical and parabolic equations and systems.

Solving nonstationary dynamic problems on base of the BIE method requires entering the notion of the *generalized solution*. That is connected with particularity of the fundamental solution of hyperbolic equations, which belongs to the class of generalized functions. Besides classical notion to differentiability of the decisions for hyperbolic equations sharply narrows the class of useful for applications problems. In particular, typical physical processes, being accompanied shock waves, are not described by differentiated solutions of such equations.

Here BIE method is elaborated for determination of nonstationary solutions of wave equations in bounden region in R^N with boundary Dirihlet or Neumann conditions. The generalized solution of BVP are constructed on the base of dynamic analogues of Green and Gauss formulas for solutions of the wave equation in space of generalized function. Their regular integral representations and singular BIE are built for $N=1,2,3$, also at presence of shock waves.

1. The generalized solutions of the wave equation.Shock waves.The multidimensional analogue of the D'Alambert's equation is considered:

$$\square_c u \equiv \Delta u - \frac{1}{c^2} \frac{\partial^2 u}{\partial t^2} = G(x, t), \quad x \in R^N, \quad t \in R^1. \quad (1)$$

G is regular function, here and hereinafter Δ is Laplace operator, $u_{,i} = \frac{\partial u}{\partial x_i}$, $\dot{u} = u_{,t} = \frac{\partial u}{\partial t}$.

It is well known [1,2] that Eq. (1) is strictly hyperbolic, class of its solutions contains functions with breakup of derivatives. The surface of the breakup F in $R^{N+1} = \{(x, \tau \equiv ct)\}$ is the characteristic surface of Eq.(1), which satisfies to the relations: $\nu_\tau^2 - \|\nu\|_N^2 = 0$, $\nu_\tau = \nu_{N+1} < 0$, where $\nu(x, \tau) = (\nu_1, \dots, \nu_N, \nu_\tau)$ is a normal vector to F , $\|\nu\|_N = \sqrt{\nu_j \nu_j}$. In R^N this is a wave front F_t , moving with the constant speed c :

$$c = -\nu_t / \|\nu\|_N, \quad \nu_t = c \nu_\tau \quad (2)$$

On reiterative indexes i, j in product everywhere their is summation from 1 to N .

There are the Adamar's conditions on the jumps at F_t :

$$[u(x, t)]_{F_t} = 0, \quad [\dot{u} + c n_i u_{,i}]_{F_t} = 0. \quad (3)$$

Here $[f(x, t)]_{F_t} = f^+(x, t) - f^-(x, t) = \lim_{\varepsilon \rightarrow +0} (f(x + \varepsilon n, t) - f(x - \varepsilon n, t))$, $x \in F_t$, $n(x, t)$ is a unit normal vector to F_t , directed aside its spreading:

$$n_i = \frac{\nu_i}{\|\nu\|_N} = \frac{\text{grad } F}{\|\text{grad } F\|}, \quad i = 1, \dots, N; \quad (4)$$

Last equality may be written only if the equation of wave front can be present as $F(x, t) = 0$, at condition of existence $\text{grad } F$.

The class of the similar solutions of the hyperbolic equations is named *shock waves*.

From second condition (3) follows

$$\dot{u}^- + cn_i u_{,i}^- = \dot{u}^+ + cn_i u_{,i}^+, \quad x \in F_t. \quad (5)$$

If before the front $u \equiv 0$, this equality gives the useful correlation on F_t : $(\text{grad } u, n) = -c^{-1} \dot{u}$.

We notice, that tangent derivatives to characteristic surface, on the strength of continuity u , are also continues. Then

$$[u_{,\tau} \gamma_\tau + u_j \gamma_j]_F = 0 \quad \text{for } \forall \gamma \in R^{N+1} : (\nu, \gamma) = 0. \quad (6)$$

In particular if $\gamma = \gamma^j = (-\nu_j, \nu_\tau \delta_1^j, \nu_\tau \delta_2^j, \dots, \nu_\tau \delta_N^j)$, where δ_i^j is Kronecker's symbol, then we have the condition of the type:

$$[-u_{,\tau} \nu_j + u_j \nu_\tau]_F = 0 \Rightarrow [\dot{u} n_j + c u_j]_{F_t} = 0, \quad j = 1, \dots, N. \quad (7)$$

Hereinafter we shall consider the functions $u(x, t)$, which are continues together with derivatives of first and second order almost everywhere with the exclusion of finite or counting number of wave fronts, on which the conditions on jumps are satisfied (2). We shall name such solutions *classical*. Let's show that they are generalized solutions of the Eq. (1).

For this let's consider the Eq.(1) on the space of generalized functions $D'(R^{N+1}) = \{\hat{f}(x, \tau)\}$, which are determined on the space of infinitely differentiable finite functions $D(R^{N+1}) = \{\varphi(x, \tau)\}$ [2]. The value $\hat{f}$ on φ , as it's accepted, is denoted as $(\hat{f}, \varphi)$. For regular function $\hat{f}$, corresponding to local integrable f , $(\hat{f}, \varphi) = \int_{R^{N+1}} f(x, \tau) \varphi(x, \tau) dV(x) d\tau$. Here everywhere $dV(x) = dx_1 \dots dx_N$.

D e f i n i t i o n . Function $\hat{f} \in D'(R^N)$ is identified as *generalized solution* of Eq. (1) if for any $\varphi \in D(R^{N+1})$ $(\square_c \hat{f}, \varphi) \equiv (\hat{f}, \square_c \varphi) = (G, \varphi)$.

L e m m a 1.1. If $u(x, t)$ is the classical solution of Eq. (1), then $\hat{u}(x, t)$ is its generalized solution.

P r o o f. If $u(x, t)$ has a finite breakup on F then [2] $\hat{u}_{,j} = u_{,j} + [u]_F \nu_j \delta_F(x, \tau)$, where first summand on the right is a classical derivative over x_j , $\|\nu\| = 1$, δ_F is *simple layer* on F :

$$([u]_F \nu_j \delta_F(x, \tau), \varphi(x, \tau)) = \int_F [u(x, \tau)]_F \nu_j(x, \tau) \varphi(x, \tau) dF(x, \tau), \quad \forall \varphi \in D(R^{N+1}).$$

Here integral on F surfaced. On the strength of continuity u outside of wavefront

$$[u]_F = \lim_{\varepsilon \rightarrow +0} (u(x + \varepsilon n, t) - u(x - \varepsilon n, t)) = u^+(x, t) - u^-(x, t) = [u]_{F_t}.$$

So, with regard for (4), we get:

$$\hat{u}_{,j} = u_{,j} + [u]_{F_t} \nu_j \delta_F(x, \tau), \quad \hat{u}_{,jj} = u_{,jj} + [u_{,j}]_{F_t} \|\nu\|_N n_j \delta_F + \partial_j \{ \|\nu\|_N [u]_{F_t} n_j \delta_F \}.$$

On virtue of Eq.(2)

$\hat{u}_{,\tau} = u_{,\tau} + [u]_{F_t} \nu_\tau \delta_F = c^{-1} u_{,t} - \|\nu\|_N [u]_{F_t} \delta_F$, $\hat{u}_{,tt} = c^{-2} u_{,tt} - c^{-1} [u_{,t}]_{F_t} \|\nu\|_N \delta_F - \{ \|\nu\|_N [u]_{F_t} \delta_F \}_\tau$.
With regard for these equalities and Adamar's conditions (1)

$$\square_c \hat{u} = \square_c u + \{ c^{-1} [u_{,t}]_{F_t} + [n_j u_{,j}]_{F_t} \} \|\nu\|_N \delta_F(x, \tau) + \\ + c^{-1} \partial_t \{ \|\nu\|_N [u]_{F_t} \delta_F(x, \tau) \} + \partial_j \{ \|\nu\|_N [u]_{F_t} n_j \delta_F(x, \tau) \} = \hat{G}(x, t),$$

since all densities of simple and double layers on F_t are equal to zero. Really, the second summand is a zero on the strength of the second condition (3) on front. But two others are null on the strength of the first one, since their action on $D(R^{N+1})$ is defined as

$$(c^{-1} \partial_t \{ \|\nu\|_N [u]_{F_t} \delta_F(x, \tau) \} + \partial_j \{ \|\nu\|_N [u]_{F_t} n_j \delta_F(x, \tau) \}, \varphi(x, t)) = \\ = - \int_F \nu_\tau [u]_F \left(\frac{\partial \varphi}{\partial n} - \varphi_{,\tau} \right) dF(x, \tau) = 0.$$

The lemma has been proved.

R e m a r k 1. From this lemma follow that condition on fronts of the shock waves easy to get, considering classical solutions of the hyperbolic equations as generalized one. It is enough to equate to zero the density, corresponding to independent singular generalized functions - analogues of simple, double and others layers, appearing under generalized differentiation of the solutions. The determination of such conditions on base of the classical methods is more labour-consuming procedure.

R e m a r k 2. Eq.(1) allows to consider the generalized solutions with derivatives breakup also on moving surface $F(x, t) = 0$, which velocity of the motion can depend on point of the front, then $v(x, t) = -F_{,t} / \|\text{grad } F\|$. On they the Adamar's conditions (3) are fulfilled with change c to $v(x, t)$. Such solutions can be generated by the right part of equation if the support of function $G(x, t)$ enlarges in the course of time in R^N .

2. Statement of nonstationary BVP. Uniqueness of the solution. Let the solution of Eq.(1) is determined in $S^- \subset R^N$, bounded by Lyapunov's surface S ([2], p. 409), $t \geq 0$.

Initial condition: by $t = 0$

$$u(x, 0) = u_0(x) \quad \text{for } x \in S^- + S, \quad u_{,t}(x, 0) = \dot{u}_0(x) \quad \text{for } x \in S^-. \quad (8)$$

We'll consider two BVP problems, consequently Dirihlet or Neumann *boundary conditions*:

$$u = u_S(x, t) \quad \text{for } x \in S, \quad t \geq 0 \quad (\text{first BVP}); \quad (9)$$

$$\frac{\partial u}{\partial n} = p(x, t) \quad \text{for } x \in S, \quad t \geq 0 \quad (\text{second BVP}). \quad (10)$$

Here we denote as $n = (n_1, \dots, n_N)$ a unit vector of external normal to S , $D^- = S^- \times R^+$, $R^+ = [0, \infty)$.

It is supposed that initial and boundary functions $u_0(x)$ and $u_S(x, t)$ are continues, $\dot{u}_0(x)$ and $p(x, t)$ piecewise continues. For the first BVP the boundary and initial conditions are coordinated:

$$u_0(x) = u_S(x, 0) \quad \text{for } x \in S. \quad (11)$$

On wave fronts, if they appear, Adamar's conditions (3) are fulfilled.

Notice that shock waves always appear if there is not coordination condition between initial velocities and velocities at boundary:

$$\dot{u}_0(x) = \dot{u}_S(x, 0) \quad \text{for } x \in S, \quad (12)$$

That is typically for physical problems. In this case at initial moment of time at the boundary D the shock wave front is formed, which spreads with the speed c in R^N . For building continuously differentiated solutions this condition is necessary. Here we don't suppose it.

It's supposed that initial conditions have been given and the one of the boundary conditions is known accordingly considered problem.

We enter the functions $E = 0, 5 \left(u_{,\tau}^2 + \sum_{j=1}^N u_{,j}^2 \right)$, $L = 0, 5 \left(u_{,\tau}^2 - \sum_{j=1}^N u_{,j}^2 \right)$.

L e m m a 2.1. *If $u(x, t)$ is a classical solution of Eq. (1) then*

$$[E]_{F_t} = -c^{-1} \left[\dot{u} \frac{\partial u}{\partial n} \right]_{F_t}, \quad [L(x, t)]_{F_t} = c^{-2} \left(\dot{u}^- + c \frac{\partial u^-}{\partial n} \right) [\dot{u}]. \quad (13)$$

P r o o f. On the strength of equality $[ab] = a^+[b] + b^-[a]$, with regard to (1) and (7), we have

$$\begin{aligned} \left[cE + \dot{u} \frac{\partial u}{\partial n} \right] &= \left[0, 5 \left(c^{-1} \dot{u}^2 + cu_{,j} u_{,j} \right) + \dot{u} \frac{\partial u}{\partial n} \right] = \\ &= 0, 5 c^{-1} \left(\dot{u}^+ \left[\dot{u} + c \frac{\partial u}{\partial n} \right] + (\dot{u}^- + cu_{,j}^- n_j) [\dot{u}] \right) + 0, 5 u_{,j}^+ [cu_{,j} + \dot{u} n_j] + \\ &\quad + 0, 5 (cu_{,j}^- + \dot{u}^- n_j) [u_{,j}] = 0, 5 c^{-1} [\dot{u}] (\dot{u}^- + cu_{,j}^- n_j) + \\ &\quad + 0, 5 [u_{,j}] (cu_{,j}^- + \dot{u}^- n_j) = 0, 5 cu_{,j}^- [u_{,j} + c^{-1} n_j \dot{u}] + 0, 5 c^{-1} \dot{u}^- [cn_j u_{,j} + \dot{u}] = 0 \end{aligned}$$

(here n is a unit normal to front F_t in R^N). Thence the first formula of the lemma follows.

Hereinafter since $[a^2] = (a^+ + a^-)[a]$, and on the strength of (3) and (6), we get second formula (13):

$$\begin{aligned} [L] &= 0, 5 \left[u_{,\tau}^2 - \sum_{j=1}^N u_{,j}^2 \right] = 0, 5 (u_{,\tau}^+ + u_{,\tau}^-) [u_{,\tau}] - 0, 5 (u_{,j}^+ + u_{,j}^-) [u_{,j}] = \\ &= 0, 5 (u_{,\tau}^+ + u_{,\tau}^-) [u_{,\tau}] + 0, 5 (u_{,j}^+ + u_{,j}^-) n_j [u_{,\tau}] = \\ &= 0, 5 \{ (u_{,\tau}^+ + n_j u_{,j}^+) + (u_{,\tau}^- + n_j u_{,j}^-) \} [u_{,\tau}] = c^{-2} \left(\dot{u}^- + c \frac{\partial u^-}{\partial n} \right) [\dot{u}]. \end{aligned}$$

R e m a r k. If before the front of the wave $u \equiv 0$, that, with use of (5), we have $[L(x, t)]_{F_t} = 0$ i.e. in this case function L continues.

T h e o r e m 2.1. *If $u(x, t)$ is classical solution of BVP, then*

$$\int_{S^-} (E(x, t) - E(x, 0)) dV(x) = - \int_0^t dt \int_{D^-} G(x, t) u_{,t} dV(x) + \int_0^t \int_S \dot{u}_S(x, t) p(x, t) dS(x) dt.$$

P r o o f. Multiplying Eq.(1) on $u_{,\tau}$ in the field of differentiability, after simple transformations we get:

$$E_{,\tau} - (u_{,\tau} u_{,j})_{,j} = -u_{,\tau} G. \quad (14)$$

But now we integrate Eq.(14) on D^- , with regard to dividing the area of integration D^- on the parts, bounded inside by wave fronts F_k .

Let us consider the left part of this equation as divergency of some vector in R^{N+1} , which is continues and differentiable in region between fronts. Then, using Ostrogradskiy-Gauss theorem in R^{N+1} , we get

$$\begin{aligned} & \int_{D^-} E_{,\tau} dV(x) d\tau - \int_{D^-} (u_{,\tau} u_{,j})_{,j} dV(x) d\tau + \int_{D^-} u_{,\tau} G(x, \tau) dV(x) d\tau = \\ & = \int_{D^-} u_{,\tau} G(x, \tau) dV(x) d\tau + \int_{S^-} (E(x, \tau) - E(x, 0)) dV(x) - \int_0^\tau \int_S (u_{,\tau} u_{,j} n_j) dS(x) d\tau + \\ & \quad + \sum_{F_k} \int_{F_k} [E \nu_\tau - u_{,\tau} u_{,j} \nu_j]_{F_k} dF_k(x, \tau) = 0, \end{aligned}$$

here $dF_k(x, \tau)$ is a differential of the area of surface in corresponding point of wave front F_k . On the strength of formulas (2) and (13), $[E \nu_\tau - u_{,\tau} u_{,j} \nu_j]_{F_k} = -c^{-1} \|\nu\|_N [cE + \dot{u} \frac{\partial u}{\partial n}] = 0$. So the last integral is equal to zero. With regard for indications for boundary function, thence the formula of the theorem follows.

R e m a r k. First condition of formulas (11) easy be get, considering Eq. (14) in $D'(R^{N+1})$: $\hat{E}_{,\tau} - (u_{,\tau} u_{,j})_{,j} = -u_{,\tau} G + \{[E] \nu_\tau - [u_{,\tau} u_{,j}] \nu_j\} \delta_F = -u_{,\tau} G - \|\nu\|_N \{[E] + [u_{,\tau} u_{,j}] n_j\} \delta_{F_t}$. Eq. (13) is fulfilled in $D'(R^{N+1})$ only if $[E + c^{-1} \dot{u} \frac{\partial u}{\partial n}]_{F_t} = 0$

T h e o r e m 2.2. *If classical solution of the first (second) BVP exists, then it is single.*

P r o o f. On the strength of linearity of the problem, it is enough to prove uniqueness of null solution. For it $G = 0$, initial and corresponding boundary conditions are null. Then, as easy see, from theorem 2.1 follows that $\int_{S^-} E(x, t) dV(x) = 0$. Since E is nonnegative, consequently $E \equiv 0 \Rightarrow u = const$. From initial conditions follows $u \equiv 0$.

T e o r e m 2.3. *If $u(x, t)$ is classical solution of the BVP, then*

$$\begin{aligned} \int_{D^-} L(x, t) dV(x) dt &= \int_{D^-} u G(x, t) dV(x) dt - \int_0^t \int_S u_S(x, t) p(x, t) dS(x) dt + \\ & \quad + c^{-2} \int_{S^-} (u \dot{u}(x, t) - u_0 \dot{u}_0(x)) dV(x) \end{aligned}$$

P r o o f. Multiplying Eq.(1) to u , after simple transformations we have:

$$L(x, \tau) + (u u_{,j})_{,j} - (u u_{,\tau})_{,\tau} = G. \quad (15)$$

Let integrate (15) over D^- with regard for its partition with wave fronts F_k . Similarly, as in

theorem 2.1, using Ostrogradskiy-Gauss theorem , we get

$$\begin{aligned} \int_{D^-} L dV(x) d\tau &= \int_{D^-} u G dV(x) d\tau + \int_{D^-} ((uu_{,\tau})_{,\tau} - (uu_{,j})_{,j}) dV(x) d\tau = \\ &= \int_{D^-} u G dV(x) d\tau + c^{-1} \int_{S^-} (uu_{,t} - u_0 \dot{u}_0) dV(x) - \int_0^\tau d\tau \int_S uu_{,j} n_j dS(x) + \\ &\quad + \sum_{F_k} \int_{F_k} [uu_{,\tau} \nu_\tau - uu_{,j} \nu_j]_{F_k} dF_k(x, \tau). \end{aligned}$$

On the strength of Adamar's conditions (3), the last summand is equal to zero. So, with regard for conditions on boundary (9), (10), we get the formula of the theorem.

Remark. Theorem 2.1. is the law of energy conservation. Similarly it's convenient to use such law with regard for conditions on front of the shock waves for proof of uniqueness of BVP solutions in nonstationary problems of mathematical physics .

3. The dynamic analogue of Green formula in $D'(R^{N+1})$. For building the BVP solution we consider it in $D'(R^{N+1})$. For this we introduce the characteristic function of semicylinder $D^- : H_D^-(x, t) \equiv H_S^-(x) H(t)$, where $H_S^-(x)$ is characteristic function of set S^- , which is equal to 0,5 on S , $H(t)$ is Heaviside function equal to 0,5 by $t = 0$. It's easy to show that

$$\frac{\partial H_D^-}{\partial x_j} = -n_j \delta_S(x) H(t), \quad \frac{\partial H_D^-}{\partial t} = -n_j H_S^-(x) \delta(t). \quad (16)$$

Hereinafter we will consider the generalized functions

$$\hat{u}(x, t) = u(x, t) H_D^-(x, t), \quad \hat{G}(x, t) = G(x, t) H_S^-(x) H(t),$$

where $u(x, t)$ is the classical solution of BVP.

Let us define the action of wave operator on $\hat{u}(x, t)$. Since $[u]_D = -u$, executing generalized differentiation and using Eq. (1) in the field of differentiability, we get

$$\begin{aligned} \square_c \hat{u}(x, t) &= -\frac{\partial u}{\partial n} \delta_S(x) H(t) - H(t) (un_j \delta_S(x))_{,j} - \\ &\quad - c^{-2} H_S^-(x) u_0(x) \dot{\delta}(t) - c^{-2} H_S^-(x) \dot{u}_0(x) \delta(t) + \hat{G}, \end{aligned} \quad (17)$$

where $\beta(x, t) \delta_S(x) H(t)$ is the simple layer on $\{S \times R^+\}$, $\delta(t)$ is Dirak function, $\frac{\partial u}{\partial n} = u_{,i} n_i$ is a derivative along normal vector n to S . Notice that density of simple and double layers are here defined by use of boundary conditions (a part from which, in depend on solved BVP, are known) and by given initial conditions.

We can present the solution of the Eq. (17) as the convolution of its right part with the fundamental solution $\hat{U}(x, t)$:

$$\square_c \hat{U} = \delta(x) \delta(t), \quad (18)$$

$$\hat{U} = 0 \quad \text{for } t < 0 \text{ and } \|x\| > ct \quad (\text{radiation conditions}). \quad (19)$$

We shall name it *Green function* of Eq. (1).

The solution (16) is represented in the manner of the following convolution (*):

$$\begin{aligned} \hat{u}(x, t) = u(x, t) H_S^-(x) H(t) = & -\hat{U}(x, t) * \frac{\partial u}{\partial n} \delta_S(x) H(t) - \left(\hat{U} * u n_j \delta_S(x) H(t) \right)_{,j} - \\ & -c^{-2} \left(\hat{U} * H_S^-(x) u_0(x) \right)_{,t} - c^{-2} \hat{U} * H_S^-(x) \dot{u}_0(x) + \hat{G} * \hat{U}, \end{aligned} \quad (20)$$

where symbol " $*$ " means that convolution on x is taken only over x . Moreover the solution (20) is single solution in the class of functions, allowing the convolution with $U(x, t)$.

If initial data and G are null then

$$\hat{u}(x, t) = -\hat{U}(x, t) * \frac{\partial u}{\partial n} \delta_S(x) H(t) - \hat{U}_{,j} * u n_j \delta_S(x) H(t). \quad (21)$$

This formula expresses the BVP solution through boundary values of function u and its normal derivative and similar to Green formula for solutions of Laplace equation [2]. However, on the strength of particularities of the fundamental solutions of hyperbolic equations on wave fronts, which type depends on space dimensionality, its integral representation gives the dispersing integrals (in second summand).

For building its regular integral presentation we introduce the antiderivatives on t functions:

$$\hat{W}(x, t) = \hat{U}(x, t) * \delta(x) H(t) = \hat{U}(x, t) *_{t} H(t) \Rightarrow \partial_t \hat{W}(x, t) = \hat{U}(x, t); \quad (22)$$

$$\hat{H}(x, m, t) = \frac{\partial \hat{W}(x, t)}{\partial x_i} m_i = \frac{\partial \hat{W}(x, t)}{\partial m}.$$

Easy to see that they also are solutions of Eq.(1) when $G = H(t)\delta(x)$ and $G = H(t)\frac{\partial \delta(x, t)}{\partial m}$ accordingly.

T h e o r e m 3.1. *In $D'(R^{N+1})$ BVP solution satisfies to the equation:*

$$\begin{aligned} \hat{u}(x, t) = & -\hat{U}(x, t) * \frac{\partial u}{\partial n} \delta_S(x) H(t) - \left(\hat{W}_{,j} * u n_j \delta_S(x) H(t) \right) - \\ & - \hat{W}_{,j} *_{x} u_0(x) n_j(x) \delta_S(x) - c^{-2} \hat{U} *_{x} H_S^-(x) \dot{u}_0(x) - c^{-2} \left(\hat{U} *_{x} H_S^-(x) u_0(x) \right)_{,t} + \hat{G} * \hat{U} y u \end{aligned} \quad (23)$$

P r o o f. It's easy to show, using determination of derivative of generalized functions and continuity u , that

$$(u n_j \delta_S(x) H(t))_{,t} = \dot{u}(x, t) n_j(x) \delta_S(x) H(t) + u(x, 0) n_j(x) \delta_S(x) \delta(t).$$

Using this relation, last formula(22) and properties of convolution differentiation, we have

$$\left(\hat{U} * u n_j \delta_S(x) H(t) \right)_{,j} = \hat{W}_{,j} * \dot{u}(x, t) n_j(x) \delta_S(x) H(t) + \hat{W}_{,j} *_{x} u_0(x) n_j(x) \delta_S(x)$$

Substituting these correlations in formula (20), we get the formula of the theorem.

From theorem follow that solution of the problem is completely defined by boundary value of the normal derivatives of $u(x, t)$ and its velocities $\dot{u}$. In analogy with presentation of the solutions of the Laplace equation, these formulas possible to name *dynamic analogue of Green formula*.

The formula (23) possesses the advantage in comparison with (20), since it allows immediately to go to its integral record without regularization of integral function on front, as it was earlier

offered in paper [3]. For $x \in S$ formula (23) gives, what we show hereinafter, boundary integral equation for BVP solution. If one of the boundary function is known, then after solving BIE on border, we can find second boundary function. Whereupon formula (23) gives the solution $u(x, t)$ in D^- .

4. Dynamic analogue of Gauss formula. We introduce the functions

$$U(x, y, t) = \hat{U}(x - y, t), \quad W(x, y, t) = \hat{W}(x - y, t), \quad H(x, y, m, t) = \hat{H}(x - y, m, t),$$

which, on the strength of properties of symmetries of the wave operator and δ -functions, satisfy to following symmetries correlations:

$$\begin{aligned} U(x, y, t) &= U(y, x, t), \quad W(x, y, t) = W(y, x, t), \quad \frac{\partial W}{\partial x_j} = -\frac{\partial W}{\partial y_j}, \\ H(x, y, m, t) &= -H(y, x, m, t) = -H(x, y, -m, t). \end{aligned} \quad (24)$$

L e m m a 4.1. *In $D'(R^{N+1})$ the dynamic analogue of Gauss formula has the form:*

$$-\hat{W}_{,i} *_{x} n_i(x) \delta(x) - c^{-2} \left(\hat{U} *_{x} H_S^-(x) \right)_{,t} = H_D^-(x, t) \quad (25)$$

P r o o f. If in both parts of Eq. (18) for U the convolution with $H_D^-(x, t)$ are taken, after using a property of convolution differentiation and formula (16) we get:

$$-U_{,j} * n_j(x) \delta_S(x) H(t) - c^{-2} (U_{,t} * H_S^-(x) \delta(t)) = H_D^-(x, t)$$

With regard for (22), flipping differentiation on t in the first summand and executing convolution on t in second one, we get the formula of this lemma. Integral record of this formula depends on dimensionality of the equation.

The formula (25) is an analogue of the known formula of Gauss for double layer potential (p.406, [2]), which gives the integral record of characteristic function of a set with use the fundamental solution of Laplace equation. The Gauss formula is often used for building the boundary integral equation of boundary value problems for elliptical equations and systems. Similarly it's possible to use the dynamic analogue of Gauss formula for constructing BIE in hyperbolic case. But here the dynamic analogue of Green formula is used for building BIE of setting problems.

Hereinafter we shall give integral representation of formulas of the theorem 3.1. and lemma 4.1 for space of dimensionality $N=1,2,3$, that is most typical for problems of mathematical physics.

5. BIE for plane BVP. When $N = 2$ we have a plane problem. At first let's consider the problem with null initial conditions.

We denote as $dS(y)$ the arc length differential on S in point y , $S_t(x) = \{y \in S : r < ct\}$, $S_t^-(x) = \{y \in S^- : r < ct\}$, $r = \|x - y\|$, $dV(y) = dy_1 dy_2$.

T h e o r e m 5.1. *By $N = 2$ function $\hat{u}(x, t)$ with null initial conditions ($u_0 = 0$, $\dot{u}_0 = 0$), has following integral representation:*

$$\hat{u} = \frac{c}{2\pi} \int_0^t d\tau \int_{S_\tau(x)} \left(\frac{\partial u(y, t - \tau)}{\partial n(y)} + \frac{\tau}{r} \frac{\partial r}{\partial n(y)} \dot{u}(y, t - \tau) \right) \frac{dS(y)}{\sqrt{c^2 \tau^2 - r^2}}. \quad (26)$$

Or, if to change the integration order, it has the form :

$$\begin{aligned}\hat{u}(x, t) = & \frac{c}{2\pi} \int_{S_t(x)} dS(y) \int_{r/c}^t \frac{\partial u(y, t - \tau)}{\partial n(y)} \frac{d\tau}{\sqrt{(c^2\tau^2 - r^2)}} + \\ & + \frac{c}{2\pi} \int_{S_t(x)} \frac{1}{r} \frac{\partial r}{\partial n(y)} dS(y) \int_{r/c}^t \frac{\tau \dot{u}(y, t - \tau) d\tau}{\sqrt{(c^2\tau^2 - r^2)}}.\end{aligned}$$

For $x \in S$ second integral on the right part is singular and taken in value principle sense.

P r o o f. For $N = 2$ Green function has a following type ([2], p. 206)

$$\hat{U}(x, t) = -\frac{cH(ct - R)}{2\pi\sqrt{(c^2\tau^2 - r^2)}}, \quad R = \sqrt{x_1^2 + x_2^2}. \quad (27)$$

Calculating from formula (22) we determine

$$\hat{W}(x, t) = -\frac{H(ct - R)}{2\pi} \ln \left(\frac{ct - \sqrt{c^2t^2 - R^2}}{R} \right), \quad \hat{H}(x, m, t) = -\frac{ct H(ct - R)}{2\pi\sqrt{c^2t^2 - R^2}} \frac{x_j m_j}{R^2}. \quad (28)$$

If to write the convolutions (23) in integral type with regard for these relations, that we get the formula of the theorem. Notice that for $x \notin S$ integrals on the right are regular and therefore for such x on the right and on the left the regular functions stand. We shall prove formula (26) also for $x \in S$ with regard for determinations $H_S^-(x)$.

We denote ε -vicinity of the point x ($\varepsilon \ll ct, t > 0$) through $\text{III}_\varepsilon(x) = \{y : r < \varepsilon\}$, $S_\varepsilon^-(x) = S^- - \text{III}_\varepsilon(x)$, $S_\varepsilon^+(x) = S^+ - \text{III}_\varepsilon(x)$, $O_\varepsilon(x) = \{y \in S : r \leq \varepsilon\}$, $S_\varepsilon = S - O_\varepsilon$, $\text{III}_\varepsilon^- = S^- \cap \text{III}_\varepsilon$, $\text{III}_\varepsilon^+ = S^+ \cap \text{III}_\varepsilon$, $\Gamma_\varepsilon^\pm(x) = \{y \in S^\pm : r = \varepsilon\}$, $r_{,j} = \frac{\partial r}{\partial y_j}$.

Let $x^* \in S$. We transform the sidebar S in vicinity of the point x^* , avoiding it on ε -semicircle in $S^-(\varepsilon \ll ct, t > 0)$. From dynamic analogue of the Green formula for sidebar $S_\varepsilon + \Gamma_\varepsilon^-$ we have in point $x = x^*$

$$\begin{aligned}0 = & \frac{c}{\pi} \int_{S_\varepsilon(x^*) + \Gamma_\varepsilon^-(x^*)} H(ct - r) dS(y) \int_{r/c}^t \frac{\partial u(y, t - \tau)}{\partial n(y)} \frac{d\tau}{\sqrt{c^2\tau^2 - r^2}} + \\ & + \frac{c}{2\pi} \int_{S_\varepsilon(x^*)} H(ct - r) \frac{1}{r} \frac{\partial r}{\partial n(y)} dS(y) \int_{r/c}^t \frac{\tau \dot{u}(y, t - \tau) d\tau}{\sqrt{c^2\tau^2 - r^2}} + \\ & + \frac{c}{2\pi} \int_{\Gamma_\varepsilon^-(x^*)} H(ct - r) \frac{1}{r} \frac{\partial r}{\partial n(y)} dS(y) \int_{r/c}^t \frac{\tau \dot{u}(y, t - \tau) d\tau}{\sqrt{c^2\tau^2 - r^2}}.\end{aligned}$$

For $\varepsilon \rightarrow +0$ first integral, on the strength of weak singularity of the integral functions, strives to integral on S , second one tends to integral in the value principle sense, which also exists, since integral function has a singularity of the type r^{-1} and contains function $\frac{\partial r}{\partial n(y)} = n_j(y) \frac{(y_j - x_j)}{r}$, which for $y \rightarrow x$ asymptotically equivalent to $\frac{\partial r}{\partial n(x)} = n_j(x) \frac{(y_j - x_j)}{r}$, antisymmetric relative to point x .

Let consider the last summand, denoted $J_\varepsilon(x)$. On Γ_ε^- there are $r = \varepsilon$, $\frac{\partial r}{\partial n(y)} = -1$, $dS(y) = \varepsilon d\theta$, where θ is an polar corner with top in point x ; θ_1 and θ_2 are the corners of endpoints Γ_ε^- , numbered in order at pass-by of the sidebar Γ_ε^- in positive direction. With regard for these relations

$$J_\varepsilon(x) = \frac{c}{2\pi} \int_{\theta_1}^{\theta_2} \frac{\varepsilon}{\varepsilon} d\theta \int_{\varepsilon/c}^t \frac{\tau \dot{u}(y, t - \tau) d\tau}{\sqrt{c^2 \tau^2 - r^2}} = \frac{(\theta_2 - \theta_1) c}{2\pi} \int_{\varepsilon/c}^t \frac{\tau \dot{u}(y, t - \tau) d\tau}{\sqrt{c^2 \tau^2 - r^2}},$$

$$\lim_{\varepsilon \rightarrow 0} (\theta_2 - \theta_1) = -\pi, \quad \lim_{\varepsilon \rightarrow 0} J_\varepsilon(x) = -0,5 \int_0^t \dot{u}(y, t - \tau) d\tau = -0,5 u(x, t).$$

Passing this summand in the left part, with regard for determinations: $H_D^-(x, t) = 0,5$ for $x \in S$, we get the formula of theorem on the border also. Since on the left and on the right in Eq. (26) the regular generalized functions stand, on the strength of Dyubua-Reymon lemma ([2], p. 97) this equality, which equitable in generalized functions class, also equitable in classical sense. This theorem has been proved.

By solving the BVP the Eq. (26) gives the boundary integral equation for determination of unknown boundary function:

$$\pi u_S(x, t) =$$

$$= \int_{S_t(x)} dS(y) \int_{r/c}^t \frac{\partial u(y, t - \tau)}{\partial n(y)} \frac{d\tau}{\sqrt{\tau^2 - r^2/c^2}} + V.P. \int_{S_t(x)} \frac{1}{r} \frac{\partial r}{\partial n(y)} dS(y) \int_{r/c}^t \frac{\tau \dot{u}_S(y, t - \tau) d\tau}{\sqrt{\tau^2 - r^2/c^2}}.$$

In the case of the first BVP the left part of this equation and the second integral on the right are known, they are defined with use of boundary condition, but the first integral contains the kernel with weak singularity on front of the Green function. Solving it we define the normal derivative of u on boundary whereupon formula (26) allows to compute the solution in any point.

In the case of the second BVP we have *singular* BIE for determination of unknown boundary values of u . After its solving we can define $u(x, t)$ on boundary whereupon formula (26) defines the solution fully.

In the case of nonzero initial conditions the following theorem gives the solution of problems.

T h e o r e m 5.2. *By $N = 2$ BVP solution has the following integral representation:*

$$2\pi \hat{u} = \int_0^t d\tau \int_{S_\tau(x)} \left(\frac{\partial u(y, t - \tau)}{\partial n(y)} + \frac{1}{r} \frac{\partial r}{\partial n(y)} \tau \dot{u}(y, t - \tau) \right) \frac{dS(y)}{\sqrt{\tau^2 - (r/c)^2}} +$$

$$+ \frac{\partial}{\partial t} \int_{S_t^-(x)} \frac{u_0(y) dV(y)}{c\sqrt{c^2 t^2 - r^2}} + \int_{S_t^-(x)} \frac{\dot{u}_0(y) dV(y)}{c\sqrt{c^2 t^2 - r^2}} + c \int_0^t d\tau \int_{S_\tau^-(x)} \frac{G(y, t - \tau) dV(y)}{\sqrt{c^2 \tau^2 - r^2}} +$$

$$- \int_{S_t(x)} u_0(y) \frac{ct}{r} \frac{\partial r}{\partial n(y)} \frac{dS(y)}{\sqrt{c^2 \tau^2 - r^2}}$$

Proof follows from theorem 4.1. and 5.2. if to write the convolution with initial data in integral type. Here the integrals from the second before the fourth ones comply with Poisson formula for

Cauchy problem . The last additional summand with initial data is conditioned presence of the boundary.

Using relations (27), dynamic analogue of Gauss formula possible to be write in integral type.

L e m m a 5.1. *By $N = 2$ dynamic analogue of Gauss formula has the following type:*

$$V.P. \int_{S_\tau(x)} \frac{1}{\sqrt{1 - (r/ct)^2}} \frac{\partial}{\partial n(y)} \left(\ln \frac{1}{r} \right) dS(y) + \frac{\partial}{c\partial t} \int_{S_t^-(x)} \frac{dV(y)}{\sqrt{c^2 t^2 - r^2}} = 2\pi H_S^-(x) H(t),$$

where integral in the value principle sense is taken for boundary points .

P r o o f. Formula of the lemma 4.1 , with regard for (27),(28), possible to write so

$$- \int_{S_t(x)} \frac{ct}{\sqrt{c^2 t^2 - r^2}} \frac{1}{r} \frac{\partial r}{\partial n(y)} dS(y) + \frac{\partial}{c\partial t} \int_{S_t^-(x)} \frac{dV(y)}{\sqrt{c^2 t^2 - r^2}} = 2\pi H_S^-(x) H(t). \quad (29)$$

From this formula by elementary transformations we get the formula of the lemma.

We shall show that this equality, equitable in the field of regularity , is saved also for $x \in S$ if the singular integral on the left, which contains the strong singularity on r , is taken in the value principle sense.

Similarly (29) for area without and with ε -vicinity of the point x we get

$$\begin{aligned} & - \int_{S_\varepsilon + \Gamma_\varepsilon^-} \frac{ct H(ct - r)}{\sqrt{c^2 \tau^2 - r^2}} \frac{1}{r} \frac{\partial r}{\partial n(y)} dS(y) + \frac{\partial}{c\partial t} \int_{S_\varepsilon^-(x)} \frac{H(ct - r) dV(y)}{\sqrt{c^2 \tau^2 - r^2}} = 0, \\ & - \int_{S_\varepsilon + \Gamma_\varepsilon^+} \frac{ct H(ct - r)}{\sqrt{c^2 \tau^2 - r^2}} \frac{1}{r} \frac{\partial r}{\partial n(y)} dS(y) + \frac{\partial}{c\partial t} \int_{S_\varepsilon^-(x) + \text{III}_\varepsilon} \frac{H(ct - r) dv(y)}{\sqrt{c^2 \tau^2 - r^2}} = 2\pi. \end{aligned} \quad (30)$$

Under $\varepsilon < ct$ integrals on $\Gamma_\varepsilon^\pm$, III_ε are easy calculated by transition to polar coordinate system. On $\Gamma_\varepsilon^\pm$ $\frac{\partial r}{\partial n(y)} = \pm 1$ consequently to the sign. Mark that

$$\begin{aligned} I_\varepsilon(x, t) &= \int_{\text{III}_\varepsilon(x)} \frac{H(ct - r) dV(y)}{\sqrt{c^2 \tau^2 - r^2}} = \int_0^{2\pi} d\theta \int_0^\varepsilon \frac{r dr}{\sqrt{c^2 \tau^2 - r^2}} = 2\pi(ct - \sqrt{c^2 \tau^2 - r^2}) \Rightarrow \\ & \lim_{\varepsilon \rightarrow 0} \frac{\partial I_\varepsilon}{\partial t} = 2\pi c \lim_{\varepsilon \rightarrow 0} \left(1 - \frac{ct}{\sqrt{c^2 \tau^2 - r^2}} \right) = 0 \text{ for } \forall t > 0. \end{aligned}$$

If both equalities in (30) to add and divide on 2, with regard for properties of the symmetries of integral function (24) and going to limit on $\varepsilon \rightarrow 0$, then

$$-V.P. \int_{S_t(x)} \frac{ct}{\sqrt{c^2 \tau^2 - r^2}} \frac{1}{r} \frac{\partial r}{\partial n(y)} dS(y) + \frac{\partial}{c\partial t} \int_{S_t^-(x)} \frac{dV(y)}{\sqrt{c^2 \tau^2 - r^2}} = \pi.$$

So, with regard for determinations $H_S^-(x)$, formula (29) is equitable for any x .

6. BIE of BVP for N=3.

T h e o r e m 6.1. *For $N = 3$ BVP solution is represented in the form:*

$$\begin{aligned} 4\pi\hat{u}(x, t) = & \int_{S_t(x)} \left\{ \frac{1}{r} \frac{\partial u(y, t - r/c)}{\partial n(y)} + \frac{\dot{u}(y, t - r/c)}{c} \frac{\partial \ln r}{\partial n(y)} \right\} dS(y) - \\ & - V.P. \int_{S_t(x)} u(y, t - r/c) \frac{\partial}{\partial n(y)} \frac{1}{r} dS(y) + c^{-1} \frac{\partial}{\partial t} \int_{S_t(x)} u_0(y) \frac{\partial \ln r}{\partial n(y)} dS(y) + \\ & + \int_{r=ct} \frac{\dot{u}_0(y)}{c^2 t} H_S^-(y) dS(y) + \frac{\partial}{\partial t} \int_{r=ct} \frac{u_0(y)}{c^2 t} H_S^-(y) dS(y) - \int_{S_t^-(x)} \frac{G(x, t - r/c)}{r} dV(y). \end{aligned}$$

P r o o f . For $N = 3$ Green function $\hat{U}(x, t)$ is the double layer on the cone $K_t = \{(x, t) : \|x\| = ct\}$:

$$\hat{U}(x, t) = -\frac{\delta(t - R/c)}{4\pi R}, \quad R = \sqrt{x_1^2 + x_2^2 + x_3^2}. \quad (31)$$

For any $\varphi(x, t) \in D(R_4)$ it defines the linear functional:

$$(U, \varphi) = -\frac{1}{4\pi} \int_{R^3} \frac{\varphi(x, \|x\|/c)}{\|x\|} dV(x)$$

After calculation on formula (22) we find

$$\hat{W}(x, t) = -\frac{H(ct - R)}{4\pi R}, \quad \hat{H}(x, m, t) = \frac{1}{4\pi} \frac{x_j m_j}{R^2} \left(c^{-1} \delta(t - R/c) + \frac{H(ct - R)}{R} \right). \quad (32)$$

For building the integral representation of dynamic analogue of Green and Gauss formulas we use the following equalities, which be get using the determination of generalized function convolution :

$$\begin{aligned} \alpha(x) \delta(t - R/c) * f(x, t) H(t) &= H(t) \int_{S_t(x)} \alpha(x - y) f(y, t - r/c) dS(y) \\ \beta(x, t) \delta_S(x) H(t) * f(x, t) H(t) &= H(t) \int_0^t d\tau \int_{S_t(x)} \beta(y, t - \tau) f(x - y, \tau) dS(y) \\ \alpha(x) \delta(t - R/c) * \beta(x, t) \delta_S(x) H(t) &= H(t) \int_{S_t(x)} \alpha(x - y) \beta(y, t - r/c) dS(y), \\ \alpha(x) \delta(t - R/c) * \gamma(x) \delta_S(x) &= \frac{\partial}{\partial t} \int_{S_t(x)} \alpha(x - y) \gamma(y) dS(y) \\ \alpha(x) \delta(t - R/c) * H_S^-(x) &= c^{-1} H(t) \int_{r=ct} \alpha(x - y) H_S^-(y) dS(y). \end{aligned}$$

We compute the convolution in Eq.(23) with regard for these correlations. First summand in this equation is

$$-\hat{U}(x, t) * \frac{\partial u}{\partial n} \delta_S(x) H(t) = \frac{H(t)}{4\pi} \int_{S_t(x)} \frac{1}{r} \frac{\partial u(y, t - r/c)}{\partial n(y)} dS(y).$$

The second summand $(-4\pi \hat{W}_{,j} * \dot{u}(x, t) n_j(x) \delta_S(x) H(t))$ is an amount of two convolutions:

$$\begin{aligned} -\frac{1}{4\pi c} \frac{x_j}{R^2} \delta(t - \|x\|/c) * \dot{u}(x, t) n_j(x) \delta_S(x) H(t) &= c^{-1} H(t) \int_{S_t(x)} \dot{u}(y, t - r/c) \frac{\partial \ln r}{\partial n(y)} dS(y), \\ -\frac{x_j}{R^3} H(ct - \|x\|) * \dot{u}(x, t) n_j(x) \delta_S(x) H(t) &= \\ = H(t) \int_{S_t(x)} u^0(y) \frac{\partial r^{-1}}{\partial n(y)} dS(y) - H(t) \int_{S_t(x)} u(y, t) \frac{\partial r^{-1}}{\partial n(y)} dS(y) \end{aligned}$$

Third summand

$$-4\pi \hat{W}_{,j} * u_0(x) n_j(x) \delta_S(x) = \frac{\partial}{\partial t} \int_{S_t(x)} \frac{\partial \ln r}{\partial n(y)} u_0(y) dS(y) - \int_{S_t(x)} \frac{\partial r^{-1}}{\partial n(y)} u_0(y) dS(y)$$

The last three summand give the known Kirhoff formula for solution of the Cauchy problem [1,2] with initial conditions. Summing these convolutions, we get the formula of the theorem.

In the formula of the theorem in the part, depending on initial data, there is one summand conditioned by presence of the boundary S . It disappears for $t > t^*(x)$, $t^*(x) = \max_{y \in S} \frac{\|x-y\|}{c}$ since $S_t(x) = S$ and integral does not depend on t .

In formula (16) for $x \notin S$, $t > 0$ all integrals exist. Its proof for $x \in S$ like to plane case. Herewith the strong singularity has the second summand on the right. In this case

$$\lim_{\varepsilon \rightarrow 0} \int_{\Gamma_\varepsilon^-} u\left(y, t - \frac{r}{c}\right) \frac{\partial}{\partial n(y)} \frac{1}{r} dS(y) = -\lim_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon^2} \int_{\Gamma_\varepsilon^-} u\left(y, t - \frac{r}{c}\right) dS(y) = -2\pi u(x, t),$$

It's clear that for $x \in S$ formula (31) saves kind if corresponding singular integral to calculate in the value principle sense and take into account the value $H_S^-(x)$ on S .

For $x \in S$ this formula gives the BIE for problems solving, it is singular for the second BVP.

Using correlations (19)-(22) the dynamic analogue of Gauss formula also possible to write in integral form.

L e m m a 6.1. *By $N=3$*

$$\int_{S_t(x)} \frac{\partial r^{-1}}{\partial n(y)} dS(y) + \frac{1}{c} \frac{\partial}{\partial t} \left\{ \int_{r=ct} \frac{H_S^-(y)}{r} dS(y) + \int_{S_t(x)} \frac{\partial \ln r}{\partial n(y)} dS(y) \right\} = 4\pi H_S^-(x) H(t)$$

When $t > t^*(x)$ thence the known Gauss formula [2] follows:

$$\int_S \frac{\partial}{\partial n(y)} \left(\frac{1}{r} \right) dS(y) = 4\pi H_S^-(x). \quad (33)$$

Proof of this formulas similarly is like to proof of the lemma 5.1. Notice that for $N = 2$ two-dimensional analogue of the Gauss formula is not followed from lemmas 5.1.

7. Solution of the BVP for Dalamber equation (N=1).

T h e o r e m 7.1. *For $N = 1$ BVP solution has the following integral presentation*

$$\begin{aligned}
2\hat{u} = & c H(ct - |x - a_2|) \int_{|x-a_2|/c}^t u_{,x}(a_2, \tau) d\tau - c H(ct - |x - a_1|) \int_{|x-a_1|/c}^t u_{,x}(a_1, \tau) d\tau + \\
& + \operatorname{sgn}(x - a_1) H(ct - |x - a_1|) u\left(a_1, t - \frac{|x - a_1|}{c}\right) - \\
& - \operatorname{sgn}(x - a_2) H(ct - |x - a_2|) u\left(a_2, t - \frac{|x - a_2|}{c}\right) + \\
& + c^{-1} \int_{a_1}^{a_2} \dot{u}_0(y) H(ct - |x - y|) dy + u_0(x + ct) H_S^-(x + ct) + u_0(x - ct) H_S^-(x - ct).
\end{aligned} \tag{34}$$

P r o o f. Denote $x_1 = x$. In this case ([2], p.206)

$$\begin{aligned}
\hat{U}(x, t) &= -\frac{c}{2} H(ct - |x|), \quad \hat{U}_{,t} = -\frac{c}{2} \delta(t - |x|/c), \\
\hat{W}(x, t) &= -\frac{c}{2} H(ct - |x|)(ct - |x|), \quad \hat{W}_{,x}(x, t) = \frac{c}{2} H(ct - |x|) \operatorname{sgn} x,
\end{aligned} \tag{35}$$

$$\operatorname{sgn} x = \begin{cases} 1, & x > 0; \\ 0, & x = 0; \\ -1, & x < 0. \end{cases} \tag{36}$$

In this case it is impossible to use the formula (23) for building of the integral analogue of the Green formula, because in it some functions are not determined.

It is possible to get the similar formula, if put u as zero outside of given interval and considering an action of wave operator on it in the class of generalized functions. We enter otherwise to use the formula (20). We increase the domain of definition $u(x, t)$ in the band in $R^2 \times R^+$: $\{a_1 \leq x_1 \leq a_2, -\infty < x_2 < \infty, t > 0\}$. Then boundary S will consist of two direct lines $x_1 = a_1, x_1 = a_2$, which external normals have coordinates $(-1, 0)$ and $(1, 0)$ accordingly, $\frac{\partial u}{\partial n} = n_1 \frac{\partial u}{\partial x_1}$ for $x \in S$, $H_S^-(x) = H(x_1 - a_1) H(a_2 - x_1)$, $n_1 \delta_S(x) = \frac{\partial H_S^-}{\partial x_1} = -\delta(x_1 - a_1) + \delta(x_1 - a_2)$. The formula (23) of the theorem 3.1 is converted to type:

$$\begin{aligned}
\hat{u} = & \hat{U}_2 * \frac{\partial u}{\partial x} H(t) (\delta(x - a_2) - \delta(x - a_1)) + \frac{\partial \hat{W}_2}{\partial x} * \dot{u}(x, t) H(t) (\delta(x - a_2) - \delta(x - a_1)) + \\
& + c^{-2} \left(\hat{U}_{2,t} * u_0(x) H_S^-(x) \right) + c^{-2} \hat{U}_2 * \hat{G}.
\end{aligned} \tag{37}$$

Here, all convolutions are taken with Green function for $N = 2$ and its antiderivative. On base of the method of the lowering on x_2 , rolling up on x_2 , since u does not depend on x_2 , we get

$$\begin{aligned}
\hat{u} = & \hat{U}(x - a_2, t) * \frac{\partial u(a_2, t)}{\partial x} H(t) - \hat{U}(x - a_1, t) * \frac{\partial u(a_1, t)}{\partial x} H(t) + \\
& + \hat{W}_{,x}(x - a_2, t) * \dot{u}(a_2, t) H(t) - \hat{W}_{,x}(x - a_1, t) * \dot{u}(a_1, t) H(t) + \hat{W}_{,x}(x - a_2, t) u_0(a_2) - \\
& - \hat{W}_{,x}(x - a_1, t) u_0(a_1) - c^{-2} \left(\hat{U} * \dot{u}_0(x) H_S^-(x) \right) - c^{-2} \hat{U}_{,t} * u_0(x) H_S^-(x) + \hat{U} * \hat{G}.
\end{aligned} \tag{38}$$

Substituting (36) into (38) and executing integration, we get formula (34).

Easy to show that it is equitable also for $x = a_1$, $x = a_2$ (with regard for (36)). For this it is enough to write formula (34) for interval $(a'_1, a_2) = (a_1 + \varepsilon, a_2)$ $((a_1, a_2 - \varepsilon))$. Supposing $x = a_1$ and $\varepsilon \rightarrow +0$ we have:

$$\begin{aligned}
0 &= \lim_{\varepsilon \rightarrow 0} c \left\{ H(ct - d) \int_{\frac{|x-a_2|}{c}}^t u_{,x}(a_2, \tau) d\tau - H(ct - \varepsilon) \int_{\frac{\varepsilon}{c}}^t u_{,x}(a_1, \tau) d\tau \right\} + \\
&+ H(ct - d) u(a_2, t - d/c) + \operatorname{sgn}(-\varepsilon) H(ct) u(a_1, t) + c^{-1} \int_{a_1+\varepsilon}^{a_2} \dot{u}_0(y) H(ct - |a_1 - y|) dy + \\
&+ u_0(a_1 + ct) H_S^-(a_1 + ct) + u_0(a_1 - ct) H_S^-(a_1 - ct) = \\
&= c \left\{ H(t - d/c) \int_{\frac{|x-a_2|}{c}}^t u_{,x}(a_2, \tau) d\tau - \int_0^t u_{,x}(a_1, \tau) d\tau \right\} + H(t - d/c) u\left(a_2, t - \frac{d}{c}\right) + \\
&- H(t) u(a_1, t) + c^{-1} \int_{a_1}^{a_2} \dot{u}_0(y) H(t - |a_1 - y|/c) dy + u_0(a_1 + ct) H_S^-(a_1 + ct) + \\
&+ u_0(a_1 - ct) H_S^-(a_1 - ct)
\end{aligned}$$

($d = |a_1 - a_2|$). Transferring the summand $-H(t) u(a_1, t)$ into the left part, with regard for values of the characteristic function on border, we get the formula of the theorem for left endpoint. By the similar way the formula is proved for $x = a_2$. As a result on the end of the interval (a_1, a_2) we have the following equations for determination of unknown boundary functions:

$$\begin{aligned}
u(a_1, t) &= cH(ct - d) \int_{d/c}^t u_{,x}(a_2, \tau) d\tau - cH(t) \int_0^t u_{,x}(a_1, \tau) d\tau + H(ct - d) u\left(a_2, t - \frac{d}{c}\right) + \\
&+ c^{-1} \int_{a_1}^{a_2} \dot{u}_0(y) H(ct - |a_1 - y|) dy + u_0(a_1 + ct) H(d - ct) H(t) \quad \text{для } x = a_1; \\
u(a_2, t) &= cH(t) \int_0^t u_{,x}(a_2, \tau) d\tau - cH(ct - d) \int_{d/c}^t u_{,x}(a_1, \tau) d\tau + H(ct - d) u\left(a_1, t - \frac{d}{c}\right) + \\
&+ c^{-1} \int_{a_1}^{a_2} \dot{u}_0(y) H(ct - |a_2 - y|) dy + u_0(a_2 - ct) H(d - ct) H(ct) \quad \text{для } x = a_2.
\end{aligned}$$

Under given $u_{,x}(a_k, t)$, $k = 1, 2$, we get two functional equations with lagging argument for determination u on boundary of the area, which can be solved incremental on time from $t = 0$. Under the known $u(a_1, t)$, $u(a_2, t)$ we have the system of two integral equations.

Conclusion. Using this method it is possible to build the similar formulas and boundary integral equations for solutions of BVP in space of greater dimension($N > 3$). It is particularly efficient in BVP for systems of the equations of mathematical physics, when the Green matrix of system can be constructed. Herewith the type of the equations is unessential , it can be also elliptical, as, for instance in problem of stationary diffraction of electromagnetic waves [5], or in problem of the elasticity theory [6], parabolic or mixed type in problem of thermoelastodynamics [7]. But particularly efficient this method for solving of the hyperbolic equations, where using classical methods more difficult, but sometimes and simply impossible (refer to, for instance, [6,8,9]).

The study of solubility of built BIEs presents the independent problems of the functional analysis since these equations do not pertain to well studied classical ones. However we notice that using the computing methods on base of the methods of boundary element with transition to discrete analogue BIE, allows effectively to build the solutions of like problems [10].

References

- [1] Petrovskiy I.S. The lectures about equations with partial derivatives. M., 1961
- [2] Vladimirov V.S. The equations of mathematical physics. M., 1978, 512 p.
- [3] Alekseeva L.A. Integral equations of boundary value problems for wave equation in $R_2 \times t$ // Differential equations, 1992,V.28,8.
- [4] Alekseeva L.A. The dynamic analogues of Green and Gauss formulas for solutions of the wave equation in $R_N \times t$ //Differential equations.1995. V.31, 11.
- [5] Alexeyeva L.A., Sautbekov S.S. The method of generalized function at solving of the stationary boundary value problems for Maxwell equations // Journal of computational mathematics and mathematical physics. 2000.T.40, 4.
- [6] Alexeyeva L.A. Boundary Element Method of Boundary Value Problems of elastodynamics by stationary running loads//Int. J. Engineering Analysis with Boundary Element. 1998, 11.
- [7] Alekseeva L.A., Kupesova B.N. The method of generalized function in boundary value problems of thermoelastodynamics // Applied mathematics and mechanics. 2001. V.65,2.
- [8] Alexeyeva L.A. The Generalized solutions of nonstationary boundary value problems for Maxwell equations // Journal of computational mathematics and mathematical physics. 2002. V.42, 1.
- [9] Alexeyeva L.A., Zakiryanova G.K. Generalized solutions of boundary value problems of dynamics of anisotropic elastic media// Journal of the Mechanical Behavior of Materials. 2004. 5.
- [10] Alexeyeva L.A., Dildabaev Sh.A., Zhanbyrbaev A.B., Zakiryanova G.K. Boundary Integral Equation Method in Two and three dimensional problems of elastodynamics// Int. J. Computational Mechanics.1996.V.18,2.

Lyudmila A. Alexeyeva

Professor, head of waves dynamics laboratory,
Institute of Mathematics of Education and Sciences Ministry,
Pushkin str. 125, Almaty, 050010, Kazakhstan,

tel. +7 3272 911624,

E-mail: alexeeva@math.kz

Alexeyeva L.A. Nonstationary boundary value problems for wave equation and their generalized solutions

Abstract. The multivariate analogue of D'alambert's equation in the space of generalized functions is considered. The method of generalized functions for the building of solutions of nonstationary boundary value problems for wave equations in spaces of different dimensions is elaborated. Dynamic analogues of Green and Gauss formulas for solutions of wave equation in the space of generalized functions are built. Their regular integral representations and singular boundary integral equations for solving the nonstationary problems are constructed for the spaces of the dimensions 1,2,3. The method of obtaining of conditions on fronts of shock waves is stated.